

I. The Winners of the Blue Planet Prize

1997

1997

Blue Planet Prize

Dr. James E. Lovelock
(U.K.)


Honorary Visiting Fellow of Green College,
Oxford University

Conservation International (CI)
(Headquartered in U.S.A.)



At the 1997 Blue Planet Prize Awards Ceremony, the opening slides conveyed a message to children to remind us of our responsibilities to future generations and raise the question of what each one of us can do to leave our beautiful planet intact and habitable.





His Imperial Highness Prince Akishino congratulates the laureates.



Their Imperial Highnesses Prince and Princess Akishino at the Congratulatory Party.



Dr. James E. Lovelock accepts the 1997 Blue Planet Prize.



Sir David John Wright, Ambassador of the United Kingdom to Japan (left), and Christopher J. LaFleur, Chargé d'Affaires ad interim of the United States to Japan (right), congratulate the laureates.



Dr. Russell A. Mittermeier, representing CI, accepts the 1997 Blue Planet Prize.



The Blue Planet Prize Commemorative Lectures.



Prior to the awards ceremony, the award recipients meet the press. From right: Dr. Mittermeier; Dr. Lovelock; Chairman Jiro Furumoto; and Osamu Shiragami, senior executive director of the Foundation.

Profile

Conservation International (CI)

History

- 1987 CI established by Peter A. Seligmann, the current chairman and CEO.
- 1989 Dr. Russell A. Mittermeier joined CI as president.
- 1990 First hotspot map published.
First RAP expedition conducted.
Tagua Initiative launched.
Debt-for-nature agreement signed with Madagascar (US\$5 million).
First Regional Priority Setting Workshop held in Amazonia.
- 1991 Debt-for-nature agreement signed with Mexico (US\$4 million).
CI's Geographic Information System (CISIG) released. This system is available in four languages and used in more than 350 institutions in 27 countries.
- 1992 CI participated in Rio de Janeiro Earth Summit.
CI held its first Asia-Pacific region Conservation Priority-Setting Workshop in Papua New Guinea.
- 1994 RAP expedition to Peru recorded six new species of butterflies, eight orchids, two beetles and other important new findings.
First Marine RAP Expedition conducted to the Indispensable Reefs region of the Solomon Islands.
- 1995 Formal RAP course launched to train local field biologists.
- 1996 Bolivia created the 1.9-million-hectare Madidi National Park, largely as a result of attention focused on the region through CI's RAP expeditions.
Internet network created to link 25 international biosphere reserves.
CI's RAP expeditions and long-term Peru Program led to establishment of 325,000-hectare Bahuaja-Sonene National Park.
- 1997 Owing to major efforts by CI and Brazilian partner-organization IESB, the Government of the State of Bahia created the 7,000-hectare Serra do Conduru State Park, doubling the area protected in this critically endangered hotspot region.
CI published *Reinventing the Well*, a volume highlighting the importance of "best practices" in the oil and gas industry.

Conservation International (CI) is a private, nonprofit organization that was established in 1987 with the goals of conserving global diversity and demonstrating that human beings are able to live harmoniously with nature. CI was founded by Peter A. Seligmann, the organization's current chairman and chief executive officer. The organization's president is Dr. Russell A. Mittermeier. Headquartered in Washington, D.C., CI is active in 24 countries with an international staff of nearly 400 experts in biology, forestry, conservation planning, marketing and

product development, economics, law and other areas. CI carries out many of its conservation programs through partnerships with international organizations that include the United Nations and World Bank, as well as with governments, research institutes, local nongovernmental organizations (NGOs), and major corporations. CI's activities include the identification of conservation priorities, areas of the world that are particularly rich in biodiversity and under imminent threat of destruction. CI uses the term "hotspots" to describe these areas. Through the use of cutting-edge biological research and its Rapid Assessment Program (RAP), CI quickly collects and analyzes data on potentially important sites.

Since its establishment, CI has set out to develop positive examples of people coexisting with natural habitats. The organization's activities, which are based on world-class scholarship and research, help developing countries pursue economic development while maintaining biological diversity through multifaceted efforts solidly grounded in economic theory, advanced scientific methods and respect for cultural differences.

Setting Priorities for Saving Life on Earth: Megadiversity Countries, Hotspots, and Wilderness Areas

Dr. Russell A. Mittermeier

Cristina G. Mittermeier

Cyril F. Kormos

July 2001

At the outset of the new millennium, our planet unfortunately faces a wide range of increasingly pressing environmental problems: from global warming and ozone layer depletion, to toxic waste disposal, persistent pesticides, acid rain and a host of other pollution issues impacting our air, water and soils. However, as critical as it is to address these problems, Conservation International believes that one environmental issue surpasses all others in terms of magnitude, urgency and long-term significance. That is the loss of our planet's biological diversity. Biological diversity is the sum total of life on Earth, the wealth of species, ecosystems, and ecological processes that, after all is said and done, makes our living planet what it is—the only place in the universe where we know with certainty that life exists. This diversity is not only essential as our living natural resource—our biological capital in the global bank—but doubly important because its loss is an irreversible process. We have, or can develop, technological solutions to combat most other environmental ills, especially the so-called “brown” issues mentioned above. In many cases, we simply lack the political will or the economic incentives to put them into place. However, once a species of plant or animal goes extinct, it is gone forever and will never be seen again. We now face not only the loss of individual species, but the loss of entire communities and ecosystems on which we, as living creatures, ultimately depend for our own survival.

With so many issues to confront, in so many different scenarios, and with so little time remaining in which to act, the challenge of conserving biodiversity may at times seem overwhelming. Given the seriousness of the threats to natural environments from expanding populations, development and other pressures, the question even emerges as to whether we can still have a significant impact on biodiversity losses, especially in the tropical regions of the world. Indeed, we have already lost well over half of the world's tropical forests, and we are potentially at the beginning of an extinction episode unlike anything seen since the loss of the dinosaurs some 65-million years ago. The only difference between this and the previous five major extinctions being that this one would be driven not by cosmic events, but by the actions

of a single species—our own.

Conservation International believes strongly that it is not too late to save biological diversity, and that we can in fact have a major impact. To be successful in this task, however, we must be very strategic. We must first recognize that biological diversity is by no means evenly distributed over the surface of our planet, and that much of it is concentrated in a relatively few biologically rich regions that are often under severe threat. We must, therefore, set clear priorities for conservation action in these regions. To be successful, we must also establish strong partnerships within the conservation community, and with other key actors, such as local communities and the private sector. In addition, we must bring to bear a level of funding and global support at least two orders of magnitude beyond what has been invested thus far.

The essential task of priority setting can be carried out in a variety of ways, but to Conservation International it has always been essential that biodiversity conservation be based first and foremost on biological data. Thus, we first seek to determine where the greatest species and ecosystem diversity is to be found, and then to focus on those areas with the highest levels of endemic species. Species that are found only in a particular place and nowhere else. Endemic species tend to be restricted in range and specialized in terms of habitat requirements. A growing body of research indicates that they are the most heavily impacted by human-induced changes to the environment, and therefore the first to be lost.

We have used three different kinds of priority setting: the *megadiversity country* concept, *threatened biodiversity hotspots* and *major tropical wilderness areas*. The megadiversity country concept recognizes that, of all the countries on Earth, only a small handful account for a major portion of life on Earth, including terrestrial, freshwater and marine life. Dr. Russell Mittermeier first developed this concept in 1988, based on a preliminary analysis of primate conservation priorities. In looking at where the majority of the world's primates were to be found, Dr. Mittermeier discovered that only four countries in the world, Brazil, Madagascar, Indonesia and the Democratic Republic of Congo (formerly Zaire), accounted for two thirds of all primate species. These countries also had the highest levels of endemism, with 100% of Madagascar's 55 different kinds of lemurs being endemic, and levels of endemism in Brazil and Indonesia being on the order of 100%. Subsequent analysis of other mammals, birds, reptiles, amphibians, plants and selected groups of insects led to the conclusion that 17 countries qualified for megadiversity status, with Brazil, Indonesia and Colombia topping the list, followed by Australia, Peru, Mexico, Madagascar, China and nine others. This study resulted in a book, entitled *Megadiversity: Earth's Biologically Richest Nations*, published in 1997 by the Mexican cement company, CEMEX, and Agrupación Sierra Madre, a Mexican non-governmental organization. The findings in this book indicated that these 17 countries by themselves accounted for more than two-thirds of all life forms and for the vast majority of tropical rainforests, coral reefs and other priority systems. The conclusion is obvious: we must focus most of our attention on the richest and most diverse countries roughly in proportion to the diversity they contain, and we cannot leave them out of any strategy, regardless of how difficult it may be to include some of them.

The second and most important of our priority-setting approaches is that of the *threatened biodiversity hotspots*. This concept was first developed in 1988 by British ecologist Dr.

Norman Myers, and was adopted and has been used by Conservation International and the MacArthur Foundation since 1989. In 1996, Conservation International carried out a detailed reanalysis of the concept to bring it up-to-date and to determine its validity and usefulness for the new millennium. This process reaffirmed the value of the hotspots concept and validated Myers's earlier list, but also substantially expanded upon it.

Among the most important new additions were quantitative criteria for hotspot countries. Endemism was considered the most important criterion, and plants, in this case *vascular plants*, which make up the vast majority of the Plant Kingdom, were chosen as the main group of organisms to determine whether or not an area qualified as a hotspot, primarily because most other terrestrial life forms depend on them to some degree. Data on plant endemism were combined with data on total plant diversity (all species found in a particular area, including both endemics and non-endemics) and complemented with data on diversity and endemism for four vertebrate groups (birds, mammals, reptiles and amphibians). Data on these groups were then used to refine our understanding of the importance of the hotspots, to set priorities and to provide rankings among the hotspots themselves. Thus, to qualify as a hotspot, an area must contain 0.5% of the global total of vascular plants (estimated at 300,000 species), or 1,500 species, as endemics.

The second major criterion for hotspot status, applied as a second layer of analysis after an area had qualified on the basis of plant endemism, was *degree of threat*. In order to be included on the list on the basis of this criterion, an area must have lost 70% or more of its original natural vegetation, leaving 30% or less in intact condition. The threat criterion enabled us to distinguish between areas with high diversity and endemism under severe threat (Madagascar and the Philippines) and those with equal or higher diversity and endemism but still largely intact, the so-called *major tropical wilderness areas* (Amazonia, the Congo forests of Central Africa and New Guinea) that are the focus of the third of our major priority-setting approaches.

In all, a total of 25 hotspots were identified for the terrestrial realm. These include five (20%) that are exclusively tropical rainforest; seven (28%) that include both tropical rainforest and tropical dry forest; another three (12%) with tropical rainforest, tropical dry forest and other non-forest elements (including desert scrubland or grassland formations); five (20%) that are Mediterranean-type ecosystems; three (12%) that consist of temperate forest and grasslands; one (4%) that is a mix of tropical dry forest, woodland savannas and open savannas; and one (4%) that is mainly an arid region. Of particular note is the emergence of the five Mediterranean-type systems as major priorities. These regions, which include the Cape Floristic Province of South Africa, Southwest Australia, Central Chile, the California Floristic Province, and by far the most diverse, the Mediterranean itself, are characterized by cool wet winters and hot, dry summers. They exhibit very high plant diversity and endemism. Indeed, what remains intact in these five areas accounts for only 0.2% of Earth's land surface, yet it is home to an astounding 26,743 endemic plants, 9% of the global total. Humans have also long impacted these Mediterranean-type systems, especially the Mediterranean basin, because they provide excellent climates for human habitation.

Together, the hotspots once occupied a land area of 17,541,969 km², or 11.8% of the

planet's land surface—roughly an area the size of Russia. Cumulatively, they have lost 88% of this area, meaning that only 2,127,908 km², or 12% of the original extent, remains intact. This represents just 1.4 % of the land surface of the planet, a relatively small area that is equivalent to the American states of Alaska and Texas combined, or about four times the area of France.

Our study indicates that this small portion of Earth's land surface has within it, as endemics, an astounding 133,199 vascular plant species, or 44.4% of all the world's plants. Representation of vertebrates in the hotspots is comparably impressive. Our results indicate that there are 9,681 endemic non-fish vertebrates in the hotspots, representing 35.4% of the global total. Level of vertebrate endemism varies, from the birds, which have 2,746 endemic species (27.6%) found only in hotspots, to the amphibians, which have a staggering 2,572 species, or 53.8% of the global total. Mammals totaled 1,406 endemic species, or 29.2%, and reptiles somewhat higher at 2,957 endemic species, or 37.8% of the global total.

The overall conclusions of the hotspots analysis are simple and obvious. If we have 44.4% of all plants and 35% of all non-fish vertebrates endemic to just 1.4% of Earth's land surface, and if this same 1.4% is under the most severe threat, then it is only logical to focus a significant portion of our attention on these areas in the next few decades, and especially over the next two to five years. To put it even more strongly, the hotspots are fundamental to preventing a mass extinction crisis in the early part of this new millennium. If we fail to act in these areas and lose that 1.4% of the land surface of the planet, we will lose, at the very least, those plants and animals that are found nowhere else than in the hotspots—even if all of efforts in other parts of the world are successful. Without decisive action in the hotspots, major extinctions are inevitable.

Our final priority-setting approach is the *major tropical wilderness areas approach*, which seeks to complement the hotspots by looking at high-biodiversity areas (again mainly in the tropics) that are at the opposite end of the threat spectrum. Whereas the hotspots consist of heavily exploited and often highly fragmented ecosystems greatly reduced in original extent (usually between 4% and 25% remaining), the major tropical wilderness areas are still largely intact (over 75% of original vegetation cover remaining) and have very low human population density (less than 5 people/km²). These wilderness areas have enormous importance as storehouses of biodiversity, as major watershed protection areas, and as controls against which we can measure the management of the more degraded hotspots. They are also often the last places where indigenous people have any hope of maintaining their traditional lifestyles. And they are likely to assume increasing recreational, aesthetic and spiritual value on an ever more overcrowded planet. Furthermore, since they are still under far less human pressure than the threatened hotspots (although the pressures on them are mounting rapidly), the "opportunity cost" of conservation is much lower in these areas. In other words, large-scale conservation set-asides can be achieved at far lower financial cost than in the areas where little remains and threats are high.

Few such wilderness areas have managed to persist in our rapidly changing world. The principal ones being in the Guayana Shield region encompassing Suriname, Guyana, French Guiana, Venezuela and adjacent parts of Brazil; in a large area of upper Amazonian Brazil, Colombia, Ecuador, Peru and Bolivia; in a substantial portion of the Congolese forest

block/Congo River Basin in Central Africa; and on most of the island of New Guinea and adjacent, smaller Melanesian islands (Solomon Islands, New Britain, New Ireland and Vanuatu). The major tropical wilderness areas will be the subject of the third book in this CEMEX series to appear in 2001.

To the question of whether or not we can still have a significant impact on biodiversity losses, Conservation International answers that, despite the significant challenges, we are optimistic. Success will depend on maintaining a very strategic focus on priority regions and building the alliances necessary to pursue conservation in these areas. However, given the strong support accorded to the concept of *threatened biodiversity hotspots* by the scientific community, in particular at a conference entitled "Defying Nature's End" sponsored by Conservation International in Pasadena in 2000, there is every indication that this strategic focus and partnership approach will be maintained. Thus, we firmly believe that by concentrating most of our efforts on these priority areas, we can turn today's conservation dreams into tomorrow's conservation realities.

Lecture

Biodiversity Conservation: A Global Challenge, A Global Priority

Dr. Russell A. Mittermeier
President, Conservation International

As we take our first steps into a new millennium, our planet and our species face some of the most severe threats that we have ever encountered. Amazingly, these are not only geopolitical, like the risk of nuclear war, the spread of terrorism, or other issues that have been preoccupying us since the 20th century, but rather environmental. Our planet currently faces many different environmental stresses, such as overpopulation in the developing world, over-consumption in the developed countries (and increasingly in the richer developing countries as well), global warming, ozone layer depletion, toxic waste disposal, erosion, air, soil and water pollution, and a number of others. Although many of these so-called “brown” environmental issues are more evident and may appear more urgent to our increasingly urban global society, we believe that there is one environmental issue that surpasses all others in terms of long-term significance, and that is loss of our planet’s biological diversity. This biodiversity, simply defined, is “the sum total of life on Earth,” that wealth of species, ecosystems and ecological processes that makes our living planet what it is; it’s our living resource base, our biological capital in the global “Bank,” and most importantly, its loss is an irreversible process.

We already have, or can develop technological “fixes” to most of our “brown” environmental problems, and sometimes lack only the economic incentive or political will to put them into place, but biodiversity loss is a different story. Once a species of animal or plant goes extinct, it is gone forever and will never be seen again, and we now face not just the loss of individual species, but the degradation and eventual loss of entire communities and ecosystems upon which we ourselves ultimately depend for our own survival.

One example that indicates mass extinction episodes may already be in progress is provided by the “declining amphibian phenomenon,” a global trend in which certain rare frogs and other amphibians with restricted ranges are going extinct, and more common species are becoming rare or appearing with a number of mutations and genetic defects. The golden toad (*Bufo periglens*) from a tiny area of cloud forest in Costa Rica is symbolic of this phenomenon, and has not been seen in the wild for the past decade. Regardless of all our other accomplishments, the one critical measure by which future generations will judge our success, or lack thereof, as a society as we start this new millennium will be whether or not we have the capacity to maintain the same healthy and diverse living planet that we were born into during the last century.

The mission of our organization, Conservation International, is *to conserve our planet’s*

biological diversity, and demonstrate that people and Nature can live harmoniously, and we have dedicated ourselves to achieving this complex objective. Some of the critical issues that we face in conserving biodiversity include the fundamental role of setting clear priorities, and the need for rapid action in the field in the remote areas where so much of our planet's biodiversity is still found.

How Much Do We Know about Life on Our Planet?

In terms of the most critical issues in biodiversity conservation, it is important to recognize that we are still amazingly ignorant as a culture about the rest of life on this planet. Scientists have thus far described only about 1.4- to 1.8-million species of plants, animals and microorganisms, yet estimates of total planetary diversity run as high as 10-million, 30-million or even one hundred million or even more species. If we look at the complex ecological interactions that exist between and among these different species, something that really is just in its infancy as a science, our ignorance is several orders of magnitude greater. We can send spaceships to the farthest reaches of our solar systems, and indeed spend more in the United States in a single space probe to Mars, with the rationale of seeking extraterrestrial life, than we have spent on the ground in the past decade doing biodiversity research. We put millions of bits of information on tiny computer chips, develop extremely sophisticated and complex information superhighways; and we have global institutions like the World Bank and the various United Nations agencies that invest hundreds of millions of dollars towards achieving the worthy, but poorly defined goal of sustainable development.

In spite of this, we still do not know, to within two orders of magnitude, how many other species of organisms share our planet with us. Trying to achieve sustainable development based on natural resources use in the face of such ignorance is rather like trying to construct one of those Martian space probes with nothing more than a few pieces of scrap metal, a hammer and a few rusty nails. Put another way, our wonderful technologically sophisticated 21st century society is in many ways still in the Dark Ages in terms of our understanding of the rest of life on Earth.

To give you an example of this ignorance from my personal perspective, we can look at nonhuman primates, whose study and survival have been cornerstones of my own quarter-century career in wildlife conservation. Although it's not that surprising, we still haven't catalogued all the tiny beetles of the tropical rainforest canopy, all the microorganisms living in our soil, or all the strange creatures far down in the deep ocean trenches, but one would think that we would at least know all our closest living relatives. Nonetheless, even with primates much remains to be learned. In 1992, I described a species of marmoset from the central Brazilian Amazon, and another just this year, again from the Amazon. Including these marmosets, 10 new monkeys have been discovered in Brazil since 1990, bringing the total number of primate species to 79. With 25 percent of all known species, Brazil has the greatest diversity of primates in the world. And my Brazilian colleagues have several more new forms that await scientific description.

Over the past decade in Madagascar, eight new species of lemur have been described, and several others have been rediscovered after not having been seen for more than a century.

So our ignorance of biodiversity is not limited to insects and microorganisms, but extends across the board to all forms of life.

We are similarly ignorant of the true values of biodiversity, the many goods and services that it already provides and that it has the potential to provide in the future, a nascent science that in some quarters is referred to as *valuation of biodiversity*. We talk vaguely about the future potential that the rainforests and the oceans have for new medicines and new agricultural products, we have some notion that the wild relatives of crop plants are important in maintaining genetic diversity and resistance to disease, and we can place economic value on a small number of forest products like timber, rubber, rattan and Brazil nuts that regularly enter into international commerce.

However, the vast majority of biodiversity use at regional, local and household levels throughout the tropics and the rest of the world remains largely unrecognized and unmeasured, and never enters into the national income accounts. The importance of forests in protecting watersheds and the enormous value of wetlands, though generally recognized, also remains largely unassessed in economic terms, although a recent study that placed a value of several trillion dollars per year on watersheds clearly indicated the magnitude of importance that living ecosystems represents to us.

The value that biodiversity and natural ecosystems provide us in terms of recreation is starting to be measured and recognized, especially in the enormous growth of ecotourism in recent years, but the subtle aesthetic and spiritual benefits they provide remain poorly understood. We are so far behind in capacity to value biodiversity that we do not even have in place the metrics by which this kind of analysis could be done. Clearly, much more is needed in the immediate future.

In trying to conserve global biodiversity and put a brake on spasms of extinction impending or already under way, two things concern us in particular. The first of these is the time frame under which we are operating and the other is the need to set priorities. Using tropical rainforests as an example, we have already lost on the order of 60% of all primary rainforest that existed at the beginning of this century; as if that were not bad enough, the situation in a number of countries is far more serious, with as much as 97% lost in certain parts of the world. At current rates of destruction in some of these areas, we have at best five to 10 years to halt these trends and save at least some representative remnants of what once existed.

Where to Begin? The Importance of Setting Priorities

Also of great importance is the need to set priorities. Biodiversity is by no means evenly distributed over the surface of the planet and certain regions and certain countries simply have far higher concentrations of life than others. Rather than trying to do everything, Conservation International decided early on in its history that it would focus its limited resources on those areas that are richest in biodiversity and also under the most severe threat.

Our principal priority-setting approach is based on a concept developed by British ecologist Norman Myers in two scientific papers published in 1988 and 1990. Myers recognized that a modest number of hotspot ecosystems covering a small total land area, most often in tropical forest areas, accounted for a high percentage of global biodiversity. CI scientists

revised this concept with Myers in 1998 to update and confirm his earlier findings. The results were published in 1999 in a book sponsored by the Mexican company CEMEX and entitled "Hotspots, Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions."

In this analysis, we found 25 areas around the world of critical importance to biodiversity conservation, based on the number of species present, the number of those species found exclusively in an ecosystem and the degree of threat they face. Their remaining natural habitats occupy only about 2% of the land surface of the planet, and yet have within them more than 50% of all terrestrial biodiversity and more than three-quarters of biodiversity at greatest risk. These hotspots, which include the Atlantic Forest region of Brazil, the Philippines, Madagascar and the Indian Ocean islands, and the Polynesia/Micronesia archipelago, among others, are the most endangered species-rich ecosystems with less than 25% of their original vegetation remaining.

CI's hotspots analysis is a powerful tool to help chart the course for biodiversity conservation because it makes a sometimes overwhelming issue become much more manageable. If a major portion of the world's plant and animal species are found in only 2% of the planet's land surface, then clearly a strong focus on targeted regions will maximize results. The hotspot approach can help guide investments through this critical era for life on Earth. By demonstrating that something can actually be accomplished, further investment can be stimulated and greater involvement encouraged by the private sector, international aid agencies and development banks.

Indeed, a new \$150-million fund designed to better safeguard the world's threatened biological hotspots in developing countries was recently launched as a joint initiative of Conservation International (CI), the World Bank and the Global Environment Facility (GEF), and has subsequently been joined by the MacArthur Foundation. The Critical Ecosystem Partnership Fund (CEPF) focuses primarily on the biodiversity hotspots, where it will try to advance the global conservation agenda on several fronts, resulting in improved management of protected areas and coordination in biodiversity corridors. Investments will support projects such as training, trans-national planning, local dialogue with extractive industries, conflict resolution, priority setting and consensus building, strengthening indigenous organizations and facilitation of partnerships between the private sector and protected areas.

At the other end of the threat spectrum, we have the last remaining major tropical wilderness areas. Like the hotspots, these are also rich in biodiversity, but they differ in that they still have more than 75% of their original natural vegetation and they have low human population densities of 5 people/km² or less. These areas are few and far between, but they are of great global importance for variety of reasons, among which is the fact that they are the last places where indigenous tribal societies are likely to be able to maintain any semblance of their traditional lifestyle. These include the upper Amazon region of South America; the southern Guianas and southern Venezuela, also in South America; the Congo Basin in Africa; and much of the island of New Guinea. Unfortunately, even these largely intact regions are now under assault, and like Conservation International and the Government of Suriname did in the case of the Central Suriname Wilderness Nature Reserve, it is important to set a precedent in protecting large blocks of undisturbed tropical wilderness.

To give you a few examples of these priority areas, I would like to provide a brief profile of the Atlantic Forest region of Brazil, one of the world's top-priority hotspots occurring within one of the two richest Megadiversity countries on Earth; the island of Madagascar, which is in its entirety both a threatened hotspot and a Megadiversity country; and the small South American country of Suriname and the island of New Guinea, which are perhaps the best examples of tropical wilderness areas.

Brazil's Atlantic Forest Region

Starting with the Atlantic Forest region of Brazil, this is a region that has been one of my two principal research interests over the past 25 years. Although we usually hear much more about the Amazon, the Atlantic Forest region is, without a doubt, the highest conservation priority in the country. It is a unique series of ecosystems that is quite distinct from the much more extensive Amazonian forests to the northwest, and once stretched almost continuously from the state of Rio Grande do Norte in northern Brazil, south as far as the Rio Grande do Sul, the southernmost Brazilian state, an area of about 1.2 million km², or about 3.2 times the size of Japan. When naturalists passed through this region in the beginning of the nineteenth century, they found some of the richest, tallest and most impressive forests on Earth, with an especially rich orchid and bromeliad flora and an abundant and diverse fauna.

However, this was the first part of Brazil to be colonized, and it developed into the agricultural and industrial center of the country. And this now has within its borders two of the three largest cities in all of South America and one of the two largest in the world, Sao Paulo and Rio de Janeiro. The results have been large-scale forest destruction, especially in the last three decades of rapid economic development, first for lumber and charcoal and then to make way for plantations, cattle, pasture and industry. These maps of the state of Sao Paulo show very well what has happened. Although development took place from the 16th century onward, the real pressure began in the 20th century and especially after 1950. Now, there is only about 3% left of what once was an expanse of forest in this important state.

Things are much the same in the rest of the region, and we estimate that at most 8% of the Atlantic Forest still stands. In some states, the remaining primary forest cover is 1% or less. Most of what has survived is in small, isolated, widely separate forest patches surrounded by developed and degraded land. Only in the mountains in Serra do Mar, especially along the Sao Paulo and Paraná coast and also some extents in Rio de Janeiro and Espirito Santo, do reasonably continuous stretches of forest remain, and even these are at risk.

As one might expect, the animals and plants native to the Atlantic Forest are not doing very well under such circumstances, and it is important to point out that the Atlantic Forest has one of the highest concentrations of terrestrial biodiversity on Earth. Indeed it has an estimated 20,000 higher plants, of which 6,000 are endemic; together with 261 mammals, 73 endemic; 620 birds, 160 endemic; and 260 amphibians, 128 endemic. These numbers make the Atlantic Forest one of the highest five hotspots on Earth, and if the region were a country, it would by itself rank among the top ten countries on Earth for biodiversity.

Unfortunately, much of this spectacular biodiversity is at great risk, and the primates are the species example. There are six genera and 24 different species and subspecies of monkeys

found in the Atlantic Forests. Studies that we have been conducting since 1979 indicate that fully 79% of these are endemic. Two groups of monkeys stand out among the rest, they are the miquis, genus *Brachyteles*, which are the largest of the South American monkeys, and the lion tamarins, genus *Leontopithecus*, which includes the famous golden lion tamarin and its three spectacular relatives, the golden-headed lion tamarin, the black lion tamarin, and the black-headed lion tamarin, the last of which was not discovered by science until 1990. These animals are representatives of two of the most critically endangered primates genera in the world, and they are now found only in forest remnants in the region. Their numbers are shockingly low, with only several hundred to a thousand golden lion tamarins, black lion tamarins and black-headed lion tamarins, less than 300 of the northern species of miqui and perhaps a thousand of the southern species of miqui remaining; only the golden-headed lion tamarin is somewhat more abundant, with an estimated population of 6,000 to 15,000, but even it is considered endangered. These unique and attractive animals have been the flagship species for this entire region. The golden lion tamarin has been the subject of a successful reintroduction program and a major international research and education program. All of these species have been used in a variety of campaigns and study programs that have helped to conserve not only themselves, but their remaining forest habitats as well. As a result, the miquis and tamarins which were virtually unknown to the Brazilian public at the beginning of the 1980s, are now so popular that they appear on the cover of phone books, on postage stamps, as themes of parades and in theater presentations, and they have even made it onto floats of Rio's famous Carnivale and into the local folklore.

Even more important, by using these species as symbols, we have managed to elevate this previously unknown and overlooked region to national and international attention, to the point that there are now many different conservation programs totaling tens of millions of dollars focusing on the work in the Atlantic Forest, the most recent of which is a 40-million-hectare corridors project that is being discussed by the World Bank and the European Union. As unbelievable as it may seem, much of this would not have happened had it not been for the great appeal and charisma of a handful of monkeys.

The Island of Madagascar

Switching continents now, let's take a trip to one of the world's most exotic locales, the island of Madagascar. And in moving to Madagascar we are not just traveling to another part of the world, we are really traveling back millions of years in evolutionary time. Madagascar is a unique evolutionary experiment, a living laboratory unlike anywhere else on Earth. Although it is located only about 400 kilometers off the East coast of Africa, Madagascar has been separated from the African mainland for a very long time, perhaps as long as 160-million years, and most of the plants and animal species occurring there have evolved in isolation and are found nowhere else in the world. The very high levels of endemism that have resulted, not just at the species level, but at the generic and family levels as well, really set Madagascar apart, and have contributed to its being both on the top of the list of Earth's Megadiversity countries and, like the Atlantic Forest of Brazil, one of the most threatened hotspots as well.

Since it is situated largely within the tropics, Madagascar also has very high species

diversity in certain groups of organisms, especially given its relatively small size; about 1.5 times the size of Japan. For instance, although it occupies only about 1.9% of the total land area of the African region, it has more orchid species than the whole of the African mainland, and indeed it is home to about 25% of all African plants. Its total vascular plant diversity is estimated at 11,000 to 12,000 species, of which an astounding 80% are endemic. Reptile and amphibian diversity is also quite high with some 300 species of reptiles, of which 274 are endemic; and 178 species of frogs, with 176 endemic and more species being discovered every year.

However, the most conspicuous and attractive group of organisms in Madagascar, and the one for which the country is best known, is the primates. But we are not talking about monkeys and apes, of which we usually think when we envision primates, but rather the lemurs. The living lemurs are a fantastic radiation of five families, 14 genera and 64 different species and subspecies, 100% of which are endemic to Madagascar, although two also occur on the nearby Comores, where they were almost certainly introduced by man. They include such animals as Madame Berthe's mouse lemur (*Microcebus berthae*), which, at 30 grams, is the smallest living primate, up to ones as large as the Indri (*Indri indri*), which is as big as a medium-sized dog, looks like a cross between a teddy bear and a giant panda, and moves by bounding from tree to tree like an arboreal kangaroo. And the lemur radiation includes some very strange species, most notably the mysterious Aye-aye (*Daubentonia madagascariensis*), which is surely one the strangest mammals on Earth and the only living representative of an entire primate family, the Daubentonidae.

To give some indication of the great responsibilities that Madagascar has, it alone is responsible for approximately 15% of all primate species, about 20% of all primate genera, and close to an incredible 30% of all primate families, making it the single highest priority area on Earth for conservation of these, our closest living relatives.

I should point out that we still know very little about Madagascar's great diversity and that a lot of basic biological inventories of the country's fauna and flora are still needed. This is especially true in the eastern rainforest region, which is the most diverse part of the country, but also the least known. Prior to my first trip to Madagascar in 1994, I had always wondered why little had been done there. It certainly couldn't be any more difficult to work there than, for instance, in remotes parts of the Amazon.

Now, a lot of people have a very romantic vision of what tropical forest research is like and how wonderful it is to be in the forest primeval. Well, it is wonderful in many ways, but I can assure you that it is not all coconut palms and white sand beaches. After a visit to the rainforest of eastern Madagascar, I discovered one of the reasons why so little had been done there: leeches. In all my experience in the tropical rainforest, including a number of other countries in Southeast Asia that have land leech populations, I had never before found them as abundant and as persistent as they are in the rainforest of Madagascar. Fortunately, there is a cure for them. A local plant called mangily is extremely effective in killing them and keeping them off your body as long as it is applied. But if there is no mangily around, the leeches can be a real problem.

Why are we so concerned about Madagascar? And why is it considered such as high pri-

ority? Well, although the human population is not that large, about 13-million people, or about 10% of the population of Japan in an area half as large again as Japan, there has been a long history of environmental degradation on the island. Our own species arrived there only about 1,500 to 2,000 years ago, with land-use practices from Southeast Asia and from Africa that were entirely inappropriate to the fragile Malagasy environment, and has had a major impact since that time. For example, a mosaic of forest and woodland savannah once covered the central plateau of Madagascar, but now there is little forest left. Erosion there is as serious as anywhere on the planet, and every year the rivers run red with eroded soil. A recent estimate by the World Bank indicates that Madagascar was losing more than 300-million dollars per year in future potential through erosion. Some 85% to 90% of Madagascar's natural habitats is already gone and the remainder is being chipped away for firewood and charcoal and converted for slash-and-burn agriculture.

Hunting is a problem as well. Although certain species are taboo to certain tribes, like the radiated tortoise among the Antandroy people of the far south and the Indri among the Betsimisaraka people of the eastern rainforest, others are subjected to serious hunting pressure. At risk are not only the lemurs, which are both trapped and hunted, but also other endemic species such as the coua, one of Madagascar's endemic birds.

Lest anyone believe that extinction spasms that we so often talk about are a figment of our imagination, he or she need only look at what has already been lost there over the past 2000 years in Madagascar. Among the species that disappeared are the elephant birds, which were the largest birds that ever lived and whose eggs weighed almost 10 kilograms; and fully nine genera and 16 species of giant lemurs, all of them the larger of the living species, representing almost 40% of known lemur genera. These include some spectacular creatures like *Megaladapis*, which resembled a large Australian Koala and grew to be the size of a calf, and *Archeoindris*, which was larger than an adult male gorilla and probably occupied a niche similar to that of the now-extinct North American ground sloth.

We have been very active in Madagascar, and have focused a lot of attention on key elements of the protected area network of the country, particularly on the Zahamena Reserve in the eastern rainforest region of the country and the Ankafarantsika Reserve in the western, dry deciduous forest. We have also been involved at a policy level. And our current program director, the honorable Leon Rajaobleina, was a former Ambassador to the United States and the former Minister of Finance. We have produced a protected area focused action plan for lemur conservation for Madagascar, a *Field Guide to the Lemurs of Madagascar*, to stimulate ecotourism, a much-needed source of foreign exchange and a great potential growth industry for this country. And in 1995, we held the first-ever biodiversity priority-setting workshop for the country, bringing together experts from Madagascar and all around the world.

Consequently, although we still have a very long way to go in Madagascar, and certainly cannot afford to be complacent, the fact is that there is much reason to be optimistic. We are now in an excellent position to change the course of conservation history in a unique country.

Suriname and Papua New Guinea

Finally, in terms of specific areas, I would like to focus your attention on two parts of the world,

Suriname and Papua New Guinea, that really represent the opposite end of the spectrum to the threatened hotspots of the Brazilian Atlantic Forest and the Island of Madagascar. Unlike the hotspots, where so much has already been lost, these two countries fall within the category of major tropical wilderness areas, meaning that, like the hotspots, they are high in biodiversity, but unlike the hotspots, most of their natural vegetation is still intact. These are areas where we still have some hope of achieving natural resource based sustainable development and where there is also some chance that indigenous tribal people will be able to maintain some semblance of their traditional lifestyles.

The first of these, Suriname, is a former Dutch colony that was known as Dutch Guiana. It achieved its independence from the Netherlands in 1975 and is the second of the three Guianas in size and population. It had one of the lowest human population densities on Earth, with only 400,000 people in an area of about 166,000 km² (44% the size of Japan). On top of this, most of the people live along the coast, and especially around the capital city of Paramaribo, with only about 5% living in the vast interior. Suriname also has the distinction of having the highest remaining forest cover of any country, with some 90% of its forest still in primary condition, making it one of the best countries for conservation of large intact tracts in this globally important biome.

Suriname's culture is unique and very different from the rest of South America. The coastal region is inhabited by Creoles of African origin: 31% Hindustanis of East Indian origin, 37% Javanese, 15% both of which came over as indentured servants at the turn of the century; Chinese (less than 2%); and Dutch and a variety of other groups of European origin (2%), including one of the oldest Jewish communities in the New World. Several native Amerindian tribes, among them the Trio, the Wayana, the Akurios, and the Bushnegroes or Maroons, inhabit the interior. The Bushnegroes are especially interesting. Numbering some 50,000 and divided into six tribes, they are the last runaway slave cultures in the Americas still living what is in effect a West African lifestyle. They are a fiercely independent group that has lived for centuries with little contact and little interference from the outside world.

On the other side of the planet is Papua New Guinea, one of the biologically and culturally most interesting nations on Earth. Another of the Megadiversity countries, it is also megadiverse in human groups; indeed, in spite of its relatively small size (only a little larger than Japan), it has the highest human cultural diversity of any country, with some 875 different languages spoken over its land area of some 475,369 km². What is more, most of these cultures are still intact. Like Suriname, Papua New Guinea has most of its forest intact, and some 97% of it is in the hands of traditional owners. This is rather like the United States or Brazil still being in the hands of the Native American Indians, and is very unusual for a nation that was colonized by European powers.

Back in 1997, when I wrote the first version of this essay, both the integrity of human culture and the natural environment in both Suriname and Papua New Guinea were under severe pressure, to the point that they too were at risk of being propelled into the hotspots category over the next decade. Largely overlooked for centuries, both of these countries were targeted by international timber conglomerates and international mining companies. The threat of timber companies was being driven largely by Malaysia and to a lesser extent Indonesia,

Taiwan and Korea, whereas mining interests were coming from a variety of different countries, especially Canada. These activities, which I have referred to as “the last great natural resource grab,” have a strong neo-colonial flavor, and, like the European colonialism of past centuries, leave little or nothing for the countries in question.

The attack by timber companies is particularly disconcerting since there is little respect for local forestry legislation. Widespread bribery and corruption, and much under-reporting and cheating on actual revenues, on top of the fact that concessions have five- to 10-year tax holidays, mean that the country receives very little in up-front benefits other than employment for the low income manual laborers.

Indeed, analyses of the economics of these operations, including one done for Suriname by the World Resource Institution and Conservation International, indicated that cash profits to the countries in question were very small and bordered on the insignificant. Furthermore, if real social and environmental costs were taken into the equation, these poor countries would have wound up in effect subsidizing the wealthy timber conglomerates by liquidating their only significant long-term assets, their natural resources. Needless to say, this form of predatory exploitation has no place at the beginning of the new millennium and should be resisted by the leadership of countries like the United States and Japan.

I am happy to report that, a year after I originally wrote this piece, the ecological tragedy that Suriname was about to agree to, turned into a conservation reality. Through a variety of efforts involving both high level contacts within the Suriname government and international media attention, we succeeded in convincing the Surinamers that there are better long-term alternatives for them, including ecotourism, bioprospecting, non-timber forest products and perhaps joint implementation of carbon-offset and low-impact timber harvesting. With this input, the Surinamers rejected requests from one Malaysian and two Indonesian companies for some 3-million hectares of timber concessions, representing more than 20% of their forest estate.

Indeed, in recognizing that Suriname’s future economic well-being will in part be based on the appropriate use and conservation of its natural resources, together with CI, the country created the Central Suriname Nature Reserve. This conservation corridor links three existing protected areas, making it one of the largest nature reserves in South America, and quite possibly the single-most pristine, tropical forested protected area on Earth. The Central Suriname Nature Reserve forms a corridor linking the three most important protected areas in central Suriname—the Raleighvallen, Tafelberg and Eilerts de Haan Gebergte Nature Reserves—and protects the entire upper Coppename River watershed, one of Suriname’s most important river systems.

The new reserve covers a diversity of rainforest ecosystems, and is home to all eight species of Suriname’s primates, more than 400 bird species, and other impressive wildlife, such as the jaguar, giant armadillo and giant river otter.

The Central Suriname Nature Reserve is the critically important centerpiece of a long-term conservation and development strategy for Suriname. It is our belief that the creation of this Nature Reserve will generate significant economic benefits for the country and international recognition of Suriname as a global environmental leader. We hope that this major step

by one visionary country will inspire other nations to make similar commitments to the protection of critical tropical wilderness areas around the world.

In Papua New Guinea, unfortunately, the situation is much more complex and the international companies much more deeply entrenched, sometimes resulting in political upheavals. The crisis in Bougainville, focusing on its copper mine, has been a major issue for the last decade, and the recent government of Sir Julius Chan was toppled earlier this year because of controversy over logging and mining concessions. At the same time, however, Malaysian companies are well established in many parts of Papua New Guinea and have converted large areas of forest.

Local landowners, many of whom were duped into signing over their land, are showing signs of resistance throughout the country and in other parts of Melanesia as well. In some instances, especially in the neighboring Solomon Islands, contracts were signed with illiterate landowners unable to read the documents they were signing, and logging activities are sometimes carried out 24 hours a day, using floodlights at night, to ensure that the maximum amount of timber can be extracted before the landowners realize what is being done to them and begin to resist.

The role of conservation international in these wilderness area countries is to inform them of the alternatives that exist to such predatory forms of exploitation, and to demonstrate to them successes based on rational biodiversity use in other tropical countries. One particularly successful effort was to take two delegations of Suriname parliamentarians to Costa Rica to meet with then Costa Rican President Jose Maria Figueres, one of the leading proponents of biodiversity conservation, and to see firsthand what was being done in terms of ecotourism and bioprospecting in this enlightened country.

Ecosystem Conservation: CI's Comprehensive Approach

These are some of the problems we face in biodiversity conservation and some of the priority areas in which our organization works. What I would like to do in the remainder of this presentation is to tell you a bit more about the tools that we employ in our conservation programs and the specific kinds of activities in which we are engaged. The mission of Conservation International is "to conserve global biodiversity, and to demonstrate that people can live harmoniously with Nature." Our comprehensive approach is simply referred to as "ecosystem conservation" and includes a heavy emphasis on local capacity building and the need to fully involve local people in all conservation endeavors.

We are particularly proud of the fact that we have a truly international organization. Although our main office is in Washington, D.C., we have 24 offices in the countries in which we work, and most of these are staffed entirely by nationals of the countries in question. We have staff members from more than 25 countries, speaking more than 30 different languages. The field-based programs of the organization rest heavily on what we refer to as our cornerstones, especially science, economics, policy and communications, and all of our programs have a protected area of some kind as their focal point.

A Sound Scientific Basis

Perhaps the most fundamental of all CI cornerstones is *science*, since we firmly believe that neither conservation nor sustainable development can be achieved without the soundest scientific underpinnings. To address this goal, CI, with a generous donation from CI's Board Members and founders of the Moore Foundation, Gordon and Betty Moore, has created the Center for Applied Biodiversity Science (CABS), a knowledge-based early warning system designed to identify critical issues confronting the conservation of biological diversity. It both anticipates destructive situations and is a preemptive force. The Center's mission is to strengthen our ability to respond rapidly, wisely and effectively to emerging threats to the Earth's biological diversity. To accomplish this, the Center mobilizes science, acts strategically, leverages partnerships and broadens public outreach. The Center identifies the most serious threats through a managed network of experts in related fields, such as natural resource economics, conservation biology, biogeography, forestry, marine biology and others. Experts are drawn from governmental agencies, international organizations, development agencies, private corporations, universities, research centers, nongovernmental organizations and museums.

Each year, the Center funds 10 to 15 outstanding, senior-level scientists from a range of backgrounds as Biodiversity Research Fellows, enabling them to focus on developing field-based solutions. If appropriate, the fellows field-test new theories, building upon Conservation International's presence in the hotspot regions.

Our other signature science program, the Rapid Assessment program (RAP), which uses the talents of some of the world's best field biologists to provide quick assessments of poorly known tropical ecosystems is overseen by CABS. RAP methodologies have been perfected by a team of superstar scientists working mainly in the Tropical Andean region of South America, the Melanesia region, Madagascar and, more recently, in the Caura River basin of Venezuela, the Okavango Delta of Botswana and Irian Jaya in Indonesia. Upcoming RAP expeditions include the Cutucu Mountains of Ecuador and the Kanuku Mountains of Guyana.

CI also works closely with the World Conservation Union, especially through its Species Survival Commission. This commission is the world's largest volunteer network of biodiversity experts, and currently is composed of some 8,000 members divided into more than 100 specialist groups. I have chaired the Primate Specialist Group since 1977, and we produce a wide variety of newsletters and journals through the SSC. These include a primate conservation journal, four different primate newsletters (for Asia, Africa, Madagascar and the Neotropical region), a Neotropical bat newsletter, an edentate newsletter, and a journal of cheilonian conservation and biology.

We have also just produced the latest version of the IUCN Red Data Books and Red Lists, the principal reference works on threatened species and a mainstay of the conservation community for some three decades. Finally, we are producing a Tropical Field Guide series aimed at providing ready-to-use information on a wide variety of tropical organisms, with a special goal of increasing interest in specialized ecotourism and "life-listing" and "watching" for everything from frogs and butterflies to turtles and monkeys.

Economics: Conservation through Enterprise Development

Also of great importance for Conservation International is *economics*. We recognize that it is essential to demonstrate that good biodiversity can also be good business, and that we can in fact come up with “win-win” solutions that conserve natural ecosystems and contribute to the national and local “bottom line” as well. To address these issues CI, with a generous donation from Ford Motor Company, has recently created the Center for Environmental Leadership and Business (CELB). Like CABS, CELB operates as a division of Conservation International and is headquartered in Washington, D.C. But it is governed by its own executive board of prestigious business and environment leaders.

The Center works in partnership with a wide range of companies and environmental organizations to promote business practices that reduce industry’s environmental effects and contribute to conservation. These practices also benefit business by cutting the costs associated with environmental impact and by enhancing a company’s reputation with communities, customers, employees, and shareholders. The result is what the Center calls a “net benefit” for both the global environment and for business. The Center concentrates on those industries with the greatest environmental effect on critical ecosystems and those with the potential to bring about positive environmental change, including agriculture and fisheries, forestry, energy and mining, travel and leisure, transportation, manufacturing and financial services. The Center provides an open forum where business leaders, environmentalists and academics can work together to create innovative solutions.

Another of Conservation International’s tools is the Conservation Enterprise Fund (CEF). This was created in 1999 with a \$1 million loan from the Small and Medium Enterprise (SME)—Global Environmental Facility program of the International Finance Corporation. CI acts as the financial intermediary to provide debt and equity financing of \$25,000–\$250,000 to small and medium-sized enterprises (\$5 million or less in assets) that are strategically important to conservation. The CEF is an enterprise development tool that enables conservation enterprises to expand their operations through financial leverage. CEF funds are directed to businesses engaged in activities such as agroforestry, ecotourism and wild harvest products.

Multifaceted Policy Program

The third of our cornerstones is *policy*, and it recognizes that you can have the best possible science and the strongest economic incentives and still fail in your conservation efforts if perverse national and international policies militate against what you are trying to accomplish. Our policy program is multifaceted, and deals with a wide range of issues. Beginning with the first ever debt-for-nature swap carried out by Bolivia in 1987, we have always paid attention to creative financing mechanisms and have attempted to stretch each conservation dollar as far as possible. Subsequent to that first debt swap, we have also carried out swaps in Mexico, Costa Rica and Madagascar, and several of these are producing returns.

More recently, we have pushed the concept of trust funds designed to meet the recurrent costs of managing protected areas. CI has structured national and regional conservation funds to provide long-term, steady disbursements consistent with the needs of field programs. Conservation funds have contributed significantly to conservation for specific regions, and also

to the building of leadership and institutions within communities.

The creation of a fund usually requires the creation of a board of directors drawn from local communities, from government agencies and from business groups. The process of administering the funds, making grants and dealing with regional and national institutions builds individual leadership and strengthens community interests. In essence, these funds act as catalysts in developing consensus between different interest groups such as local communities and donor agencies. Consequently, national and regional conservation funds have now become important actors in delivering resources to conservation projects.

CI's policy department is also building on our combined experiences in conservation biology, economics and policy to promote "best practices" in natural resource extraction with the goal of influencing the resource-extraction industries to become forces for biodiversity conservation, rather than threats to it.

Finally, we have worked on influencing a variety of legislative processes relating to biodiversity, from the United States to Brazil, to Madagascar, the International Finance Corporation (IFC), and the Global Environment Facility (GEF), among others.

International Communications Program

Finally, the *communications* cornerstone, which recognizes the fact that support for conservation must ultimately come from all sectors of society, from the highest-level decision makers to the poor villager living in the vicinity of a globally important protected area. Our International Communications department helps to develop regionally appropriate strategies for getting the word out and provides video and other materials ready for use in-country. Our Washington, D.C., Media department concentrates on getting press, television and Internet coverage for biodiversity, and on keeping the issue on the agenda of the national and international media at all times.

These different conservation "tools" are mixed together in a variety of ways to come up with the most effective and culturally sensitive conservation programs in each of the 23 countries in which we work, and we feel that our programs have had significant impact everywhere. Indeed, our basic philosophy is that we should not be involved in a particular area if our presence is not going to make a difference, and that we should work only where we have a real chance of "changing the course of conservation history."

Realizing the Dream of Biodiversity Conservation

After all is said and done, where do we really stand at this critical juncture in human history, as we take our first steps into this new millennium. Will the whole world wind up looking like a devastated central plateau region of Madagascar or is there a chance of developing solutions and achieving some level of harmony between Nature and the needs and aspirations of our own species?

Well, I am an optimist by nature, and I believe that we have made very significant progress, especially in the last 10 years. Ever since the Earth summit in Rio de Janeiro in June 1992, a much criticized and underrated event that I believe was the turning point in biodiversity conservation, we have seen some very important seed changes in attitude on the part of

governments, multilateral banks, bilateral aid agencies, the different branches of the United Nations and other major players.

The World Bank and other major international lending agencies, which were considered to be the enemy by the environmental community as recently as the mid 1980s, have done a major turn around and they are slowly becoming a positive force for conservation (although to be sure much needs to be done and some components of the World Bank family, especially the International Finance Corporation (IFC) lag far behind).

I had the privilege of chairing the first ever World Bank task force on biodiversity in 1998, which was actually responsible for first introducing the term "biodiversity" into the Bank lexicon. Virtually nothing was being done on the issue at that time. Now, just 13 years later, the World Bank has become a major player in biodiversity conservation, and we expect it to advance even further under the able leadership of Bank President Jim Wolfensohn.

The Global Environment Facility, a joint program of the World Bank and the United Nations Development Programme (UNDP) and the United Nations Environment Program (UNEP), and housed at the Bank, has put several hundred million dollars into biodiversity conservation since 1993. Another product of the Earth Summit, the Biodiversity Convention, provides the international legal framework and a forum for discussing biodiversity issues on a legal basis.

However, the most encouraging new development is, without a doubt, the increasing interest demonstrated by the private sector. Over the past few years, and again especially since the Earth Summit in Rio, we have seen the emergence of a new generation of corporate executives who are truly interested in doing something positive for the global environment and also understand that being "green" is good for the "bottom-line."

Here in Japan, we at CI have had an excellent relationship with the Keidanren for just the past ten years, and they have established an important nature conservation fund to support worthy projects in tropical countries. In the United States, real leadership on biodiversity conservation is coming from corporations like INTEL, United Airlines, Mobil, McDonalds, Ford Motor Company, Disney, Starbucks and a number of others, and in the tropical countries we are seeing the same trend with major participation from companies like Pulsar and CEMEX in Mexico and the Unibanco and Banco Real in Brazil. Since so much of what happens in the world is driven by or at least heavily influenced by the corporate sector, we are particularly pleased by this encouraging new trend.

The role of Japan is especially critical. As one of the world's economic powers and a major trendsetter in the Asia-Pacific region, and indeed the entire world, Japan's example is followed very closely. Japan needs to take a leadership role in environmental issues in general and in biodiversity conservation in particular, and there are very encouraging signs that this is taking place. The creation ten years ago of the very important Blue Planet Prize, which I truly believe to be the environmental equivalent to the Nobel Prize, is one such sign, and we at CI feel honored and privileged to have received this prestigious award this year.

The final message I would like to leave you with is that we all have to be upbeat and optimistic about biodiversity conservation. In spite of all the gloom and doom and endless reports of destruction, species loss, and environmental disasters, there are real success stories

out there. We have to learn from them, build on them and replicate them wherever possible. It is going to take ever increasing interest, financial support and technical support from developed countries like Japan and the United States, but I think that by working together with our colleagues from biodiversity-rich tropical countries, we can achieve solutions that will help us to maintain life on Earth for future generations and attend to real needs of people at the same time. We simply have to approach conservation as the art of the possible, and if we can do that, I see no reason why we can't turn today's conservation dreams into tomorrow's conservation realities.

Major Publications

Conservation International

President's Office

Kormos, Cyril F., B. A. Grosko, and R. A. Mittermeier. "United States Participation in International Environmental Law and Policy." *Georgetown International Environmental Law Review*, 13 (2001).

Center For Applied Biodiversity Science

da Fonseca, G.A.B., A. Balmford, C. Bibby, R.A. Mittermeier, et al. "Following Africa's Lead in Setting Priorities." *Nature*, 405 (2000), 393–394.

Brooks, T., M. Bakarr, G. Fonseca & A. Rylands. "Pessimism from Manu." *Conservation Biology*. In press.

Brooks, T. "Living on the Edge." *Nature*, 403 (2000), 26–29.

—, Ed. "Extinct Species in BirdLife." *Threatened Birds of the World*. Cambridge, U.K.: BirdLife International, 2000, 695–703.

Rice, Richard, Cheri Sugal and Gustavo Fonseca. "Sustainable Forest Management: A Review of Current Conventional Wisdom." Center for Applied Biodiversity Science, Conservation International. 2001.

Gullison, R.E., R.E. Rice, and A.G. Blundell. "Marketing Species Conservation." *Nature*, April 2000.

Brunner, A.G., R.E. Gullison, R.E. Rice and G.A.B. Fonseca. "Effectiveness of Parks in Protecting Tropical Biodiversity." *01 Science*.

Bakarr, M.I., G. Fonseca, R.A. Mittermeier, A. Rylands, and K.W. Painemilla, ed. "Hunting and Bushmeat Utilization in the African Rain Forest; Perspectives toward a Blueprint for Conservation Action." *Advances in Applied Biodiversity Science*, 2 (2001). Washington, D.C.: Center for Applied Biodiversity Science, Conservation International.

Cyril, Kormos and Layla Hughes. "Regulating Genetically Modified Organisms, Striking a Balance between Progress and Safety." In *Advances in Biodiversity Science*, 1 (2000). Washington, D.C.: Center for Applied Biodiversity Science, Conservation International.

van Roosmalen, Marc G. M., Tomas van Roosmalen, R. A. Mittermeier, and Anthony B. Rylands. "Two New Species of Marmoset, Genus *Callithrix* Erxleben, 1777 (Callitrichidae, Primates) from the Tapajos/Madeiras Interfluvium, South Central Amazon, Brazil." *Neotropical Primates*, 8 (1), March (2000).

Rylands, Anthony B., Horacio Schneider, Alfredo Langguth, R. A. Mittermeier, Colin P. Groves, Ernesto Rodriguez-Luna. "Assessment of the Diversity of New World Primates." *Neotropical Primates*, 8 (2), June (2000).

Maps

Hotspots: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions

Biodiversity Hotspots: State of the Hotspots, 2000.

Sistema De Areas Protegidas De La Amazonia; Red Latinoamericana de Cooperacion Tecnica en Parques Nacionales, otras Areas Protegidas, Flora y Fauna Silvestres- Subred de Areas Protegidas del Amazonas (SURAPA).

Vegetación de la Selva Maya (Spanish and English), Regional Analysis Program. May 2000.

Field Support Division

Ecotourism

Ecotourism Publications

Role Models for Sustainable Tourism. Ecotourism Department. September 2000.

Hillel, Oliver, Heloisa de Oliveira, Maria Emília Assy and Jamie Sweeting. *Ecotourism Capacity Building Workshops – Investing in People to Preserve the Environment*. (Original title in Portuguese: *Oficinas de Capacitação em Ecoturismo – Investindo em Pessoas para Conservar o Meio Ambiente*). Conservation International, Secretaria de Coordenação da Amazônia/Ministério do Meio Ambiente, 2000.

Edwards S., C. Holtz and O. Hillel. *Integrating Biodiversity Considerations into the Tourism Sector*. May 2000.

Sabrina Bini S., E. Assy, G. Ryan, S. Frazee, S. Edwards, O. Hillel. *Ecotourism Education and Awareness: A Manual for CI Community Extensionists*. January 2000.

Ecotourism Fact Sheets

2000. (Updated every 4—6 months).

Ecotourism Department Initiatives Paper.

Ecotourism.

Ecotourism Development Program.

Ecotourism Enterprise Development & Marketing.

Ixcán Biological Station.

Mayan Trails.

Chalalán Ecolodge.

Ghana's Central Region.

Togean Ecotourism Network Una Ecopark.

Ecotourism Brochures

2000. (Updated periodically).

Pantanal – Fazenda Rio Negro.

Scarlet Macaw Trail.

Estación Ixcán.

Eco-Escuela de Español.

Escuela Ecológico-Cultural de Idioma Español (Bio-Itzá).

Togean Ecotourism Network.

Una Ecopark.

Ecotourism Manuals

FY2001.

Feasibility Studies for Ecotourism Businesses.

Business Planning for Ecotourism Businesses.

Marketing Planning for Ecotourism Businesses.

Financial Planning for Ecotourism Businesses.

Using Marketing and Promotional Tools for Ecotourism Businesses.

Integrating Market Preferences into the Ecotourism Product Workshop.

Environmental and Cultural Interpretation Workshops for Ecotourism Enterprises.

Conservation Enterprise

Conservation Enterprise Fund Fact Sheets

Conservation Coffee. (Updated 9/2000).

Conservation Enterprise Fund. (01/2001).

Conservation Cocoa. (09/2000).

CEP Publications

Millard, E. *Enterprise as a Conservation Strategy*. Conservation Enterprise, 2000.

Morris, J. *Fostering Strategic Relationships between Community-Based Enterprises and Niche Markets*. Conservation Enterprise, 2001.